

Doctor's Order Sheet

FOLFIRI Regimen + Bevacizumab: Irinotecan - Fluorouracil -Leucovorin - Bevacizumab (Part I)

Adult Chemotherapy- Medical Oncology Metastatic Colorectal Carcinoma



Name:		
HCN:		
Date of Birth:		

Allergies:			☐ No Kno	own
Date:of Cycle Duration	: 14 days	Planned Administration Date of previous cycle: _	DD/LICATILICA A A A	
MAY PROCEED WITH DOSES AS WRITTEN IF	:			
 ANC greater than or equal to 1.5 X 10⁹/ BP less than or equal to 160/100, other LFT's and Bilirubin assessed. 	•	-	o 75 X 10º/L,	
Dipstick Urine or laboratory urinalysis for numbered cycle. If results are 2+ or 3+ of for protein, collect 24 hour urine for total	or greater tha	n or equal to 1 g/L laborato	ory urinalysis	
If this result is abnormal, dose reductions	•		oie.	
PREMEDICATIONS:				
☐ Ondansetron 16 mg PO				
☐ Dexamethasone 8 mg PO				
☐ Other:				
Authorized Prescriber:	Date	o:	Time:	
Authorized Prescriber's Signature:				
Nurse's Name:	_ Date:	DD/MONTH/YYYY	Time:	
Nurse's Signature:				

THIS IS A CONTROLLED DOCUMENT; PLEASE ENSURE THAT YOU ARE READING THE MOST RECENT VERSION. USER WILL BE SOLELY RESPONSIBLE FOR VERIFYING ITS CURRENCY AND ACCURACY.

Page 1 of 2 CP-0027 2018/06



Doctor's Order Sheet

Name:

HCN:

Date of Birth:

FOLFIRI Regimen + Bevacizumab: Irinotecan - Fluorouracil -Leucovorin - Bevacizumab

(Part II)

Adult Chemotherapy- Medical Oncology Metastatic Colorectal Carcinoma



Weight:	kg	Height:	cm	Body Surface Area	(BSA) =			
CHEMOTHERAPY (FOR HOSPITAL PHARMACY)								
☐ Irinotecan 180 mg/m² X BSA = mg IV in 500 mL D5W over 90 minutes on day 1								
☐ Dose n	nodification: 180	mg/m² X BSA -	% =	mg IV in 500 r	mL D5W			
over 90 m	inutes on day 1							
☐ Leucovorir	Leucovorin 400 mg/m² X BSA=mg IV in 250 mL D5W over 90 minutes on day 1							
Fluorouracil 400 mg/m² X BSA=mg IV push on day 1								
☐ Dose n	nodification: 400	mg/m² X BSA -	% =	mg IV push or	n day 1, THEN			
☐ Fluorouracil 2400 mg/m² X BSA=mg in D5W by continuous IV over 46 hours								
☐ Dose n	nodification: 240	00 mg/m² X BSA	% =	mg in D5W				
by continu	ous IV over 46	hours						
☐ Bevacizum	nab 5 mg/kg X v	veight (kg) =	mg IV in 100	0 mL normal saline ove	er:			
— 10 minutes during Cycle 1; — If tolerated without reaction- 10 minutes all other cycles								
(Prior to and post administration, flush lines with normal saline as Bevacizumab is not compatible with D5W; blood pressure measurement pre and post dose for first 3 cycles; and prior to Bevacizumab for subsequent cycles)								
HYDRATION/	HYDRATION/SUPPORTIVE CARE							
Atropine 0.4 mg intravenous prn for early diarrhea, abdominal cramps, rhinitis, lacrimation, diaphoresis, or flushing.								
Authorized Pres	scriber:		Date:	DD/MONTH/YYYY	Time:			
Authorized Pres	scriber's Signatu	ıre:		ID #:				
Nurse's Name:			Date:	DD/MONTH/YYYY	Time:			
Nurse's Signatu	ıre:							

Page 2 of 2 CP-0027 2018/06

THIS IS A CONTROLLED DOCUMENT; PLEASE ENSURE THAT YOU ARE READING THE MOST RECENT VERSION. USER

WILL BE SOLELY RESPONSIBLE FOR VERIFYING ITS CURRENCY AND ACCURACY.